Influenza Update: March 7, 2020

During the week ending March 7, 2020:

- There were no new influenza-related deaths and no new outbreaks reported this week. Thirty deaths and 19 outbreaks at long-term care facilities have been reported this season (since 9/29/2019).
- Based on data from King County laboratories, influenza was the most commonly identified respiratory pathogen, followed by rhinovirus and RSV. The percent of positive tests for respiratory viral pathogens was at or below rates observed this time of year and below peak levels observed during the previous five seasons.
- During the week ending March 7th, the percent of emergency department (ED) visits for influenza-like illness (ILI) was at or above baseline levels among all ages combined, and at or above peak levels observed during three of the previous five influenza seasons. The percent of ED ILI visits was on an upward trend for all age groups except for children <2 years; this may be associated with changes in careseeking behavior related to COVID-19. Among every age group except adults ages 65 years and older, the percent of ED ILI visits overall this season is higher than observed during each of the previous 5 influenza seasons. The percent of ED ILI visits has been highest among pediatric age groups, peaking at or above four of the previous five influenza seasons. This season, the percent of admissions for influenza has been highest among adults ages 65 years and older, but below levels observed during each of the previous five influenza seasons.


At a glance Week Ending 5-Year Average to Since 09/29/2019 03/07/2020 **Date** Laboratory-confirmed influenza deaths 0 30 32 0 19 46 Respiratory disease outbreaks at long-term care facilities (LTCFs) 19.1% Percentage positive influenza tests by PCR1 Season Peak 25.1% Number of labs reporting 5 Weekly Average 2244 Number of specimens tested Weekly Average 1521 Percentage of emergency department (ED) visits for ILI2 6.65% Season Peak 6.75% 5-Year Average to Date


²Based on Public Health - Seattle & King County's syndromic surveillance data representing aggregate percent of visits to King County EDs

Submissions to NREVSS by King County labs, PCR testing only											
Week#	Week ending	# Labs reporting	A (H1N1)	A (H3N2)	A (Not typed)	В	# Tested	% Flu positive			
7	2/15/2020	7	21	0	269	97	1994	19.4%			
8	2/22/2020	6	13	0	271	80	1734	21%			
9	2/29/2020	5	26	0	332	62	1761	23.9%			
10	3/7/2020	5	46	3	320	60	2244	19.1%			

Influenza results by subtype, PCR testing only (NREVSS)









¹Based on King County hospital laboratory and sentinel provider submissions to CDC's National Respiratory and Enteric Virus Surveillance System (NREVSS). NREVSS data not available for all previous seasons due to change in reporting procedures. Changes in facilities reporting to NREVSS may impact counts.


Public Health - Seattle & King County

Summary of Influenza Syndromic and Laboratory Surveillance





Note: The change from ICD-9 to ICD-10 codes in October 2015 may impact trends. Last updated Mar 8, 2020 ; 'current week' is week ending Mar 7, 2020 Baseline: Mean % ILI during non-flu weeks for previous three seasons, adding two standard deviations A non-flu week is a period of 2+ consecutive weeks where each one accounted for <2% of the season's total number of specimens that tested positive for influenza by PCR.



ALLHOSPITALS, Last Updated Mar 11, 2020

National data from CDC



A Weekly Influenza Surveillance Report Prepared by the Influenza Division Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet 2019-20 Influenza Season Week 9 ending Feb 29, 2020



A COC

"This map uses the proportion of outpatient visits to healthcare providers for influenza-like lilness to measure the ILI activity level within a state. It does not, however, measure the extent of geographic spread of flu within a state. Therefore, outbreaks occurring in a single city could cause the state to display high activity levels.

"Data collected in ILINet may disproportionately represent certain populations within a state, and therefore may not accurately depict the full picture of influenza activity for the whole state.

"Data displayed in this map are based on data collected in ILINet, whereas the State and Territorial flu activity map are based on reports from state and territorial epidemiologists. The data presented to CDC and state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.



Please report any of the following:

· Laboratory-Confirmed influenza-associated deaths

• Patients with novel or unsubtypable influenza viruses

· Outbreaks of influenza-like illness in a long-term care facility Immediately

Additional Resources: Additional King County Flu Information, Resources, and Surveillance UW Virology Laboratory Respiratory Virus Surveillance
Washington State Influenza Surveillance Update

National Influenza Update Global Influenza Update

Reporting Timeframe

Within 3 business days

Immediately

Contact Information

(206) 296-4774 Phone: (206) 296-4803



Report updated on 3/12/2020

Public Health - Seattle & King County Summary of Influenza Deaths and Long-Term Care Facility (LTCF) Influenza Outbreaks

Confirmed cases as of week 10 (ending 03/07/20)															
	201	9-2020	2018	3-2019	2017	7-2018	2016	5-2017	2015	-2016	2014	-2015	5-yea	ar avg	
Influenza Deaths in Week 10		0		3		0		4		1		0	1.6		
Influenza deaths, season to date (since 9/29/2019)		30		14		29		77		8		32		32.0	
LTCF Outbreaks in Week 10		0		2		6		2		1		1		2.4	
LTCF Outbreaks, season to date (since 9/29/2019)		19		24		48		90		11		59		46.4	
, , , , ,															
	201	019-2020 201		2018-2019		2017-2018		2016-2017		2015-2016		2014-2015		5-year avg	
Total Seasonal LTCF Outbreaks		19		43		68		92		18		65		57.2	
Flu type:															
A	10	53%	37	86%	15	22%	62	67%	7	39%	49	75%	34	59%	
В	4	21%	0	0%	6	9%	3	3%	7	39%	4	6%	4	7%	
A and B		11%	1	2%	5	7%	4	4%	0	0%	2	3%	2.4	4%	
Info not available		16%	5	12%	42	62%	23	25%	4	22%	10	15%	16.8	29%	
	201	0.2020	2016	2010	2017	7 2010	2016	2017	2015	2016	201/	1 201E	E vo	25 21/4	
		2019-2020		2018-2019		2017-2018		2016-2017		2015-2016		2014-2015		5-year avg	
Total Seasonal Influenza Deaths		30		52		50		84		16		43		49	
Flu type:															
A	18	60%	48	92%	33	66%	75	89%	10	63%	40	93%	41.2	84%	
H1/H1N1	7	23%	11	21%	1	2%	1	1%	6	38%	0	0%	3.8	8%	
Н3	1	3%	5	10%	6	12%	18	21%	1	6%	7	16%	7.4	15%	
A (not typed)	10	33%	32	62%	26	52%	56	67%	3	19%	33	77%	30	61%	
В	12	40%	2	4%	11	22%	7	8%	6	38%	3	7%	5.8	12%	
Not typed	0	0%	2	4%	6	12%	1	1%	0	0%	0	0%	1.8	4%	
Sex:															
Male	17	57%	27	52%	17	34%	41	49%	7	44%	17	40%	21.8	44%	
Female	12	40%	25	48%	33	66%	43	51%	9	56%	26	60%	27.2	56%	
Age:															
Under 5 years	1	3%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	
5 - 17	1	3%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	
18 - 44	4	13%	1	2%	0	0%	1	1%	3	19%	1	2%	1.2	2%	
45 - 64	4	13%	13	25%	7	14%	5	6%	5	31%	6	14%	7.2	15%	
65+ years		67%	38	73%	43	86%	78	93%	8	50%	36	84%	40.6	83%	
Average		64.7		73.6		81.1		81.9		64.9		81.7		76.6	
Race:															
White	17	57%	35	67%	33	66%	54	64%	12	75%	35	81%	33.8	69%	
Asian		7%	5	10%	2	4%	13	15%	2	13%	1	2%	4.6	9%	
Black	1	3%	1	2%	3	6%	4	5%	2	13%	5	12%	3	6%	
Amer Indian	1	3%	1	2%	0	0%	0	0%	0	0%	0	0%	0.2	0%	
Hispanic/Latino	3	10%	2	4%	2	4%	3	4%	0	0%	1	2%	1.6	3%	
Other	1	3%	0	0%	1	2%	1	1%	0	0%	1	2%	0.6	1%	
Unknown	5	17%	8	15%	9	18%	9	11%	0	0%	0	0%	5.2	11%	
Flu vaccine status															
Up to date	11	37%	16	31%	26	52%	39	46%	6	38%	21	49%	21.6	44%	
Not up to date	15	50%	19	37%	10	20%	20	24%	8	50%	5	12%	12.4	25%	
Unknown	4	13%	17	33%	14	28%	25	30%	2	13%	17	40%	15	31%	