Influenza Update: April 25, 2020

During the week ending April 25, 2020:

- There were no new influenza-related deaths and one new outbreak reported this week. Thirty-three deaths and 25 outbreaks at long-term care facilities have been reported this season (since 9/29/2019).
- Based on data from King County laboratories, rhinovirus was the most commonly identified respiratory pathogen, followed by adenovirus. The percent of positive tests for respiratory viral pathogens was below levels observed this time of year and below baseline levels observed during each of the previous five seasons. COVID-19 testing is not currently included in laboratory reporting of respiratory pathogens.
- During the week ending April 25th, the percent of emergency department (ED) visits for influenza-like illness (ILI) was at or below baseline levels among all ages combined, and among each individual age group. ED ILI visits have been on a downward trend over the past six weeks following a peak in week 10. Among every age group except for adults ages 65 years and older, the percent of ED ILI visits overall this season is higher than observed during each of the previous five influenza seasons. The percent of ED ILI visits has been highest among pediatric age groups, peaking at or above four of the previous five influenza seasons. This season, the percent of admissions for influenza has been highest among adults ages 65 years and older, but below levels observed during each of the previous five influenza seasons.

At a glance			
	Week Ending 04/25/2020	Since 09/29/2019	5-Year Average to Date
Laboratory-confirmed influenza deaths	0	33	45.8
Respiratory disease outbreaks at long-term care facilities (LTCFs)	1	25	56
Percentage positive influenza tests by PCR ¹	0%	Season Peak 25.10	%
Number of labs reporting	6	Weekly Average 8	
Number of specimens tested	409	Weekly Average 1481	t
Percentage of emergency department (ED) visits for ILI ²	0.94%	Season Peak 6.930	% 5-Year Average to Date 3.16%


 $^{^1}Based \ on \ King \ County \ hospital \ laboratory \ and \ sentinel \ provider \ submissions \ to \ CDC's \ National \ Respiratory \ and \ Enteric \ Virus \ Surveillance \ System \ (NREVSS)$


 $NREVSS\ data\ not\ available\ for\ all\ previous\ seasons\ due\ to\ change\ in\ reporting\ procedures.\ Changes\ in\ facilities\ reporting\ to\ NREVSS\ may\ impact\ counts\ procedures\ for\ p$

²Based on Public Health - Seattle & King County's syndromic surveillance data representing aggregate percent of visits to King County EDs.

Submissi	Submissions to NREVSS by King County labs, PCR testing only									
Week#	Week ending	# Labs reporting	A (H1N1)	A (H3N2)	A (Not typed)	В	# Tested	% Flu positive		
14	4/4/2020	6	0	0	3	1	778	0.5%		
15	4/11/2020	6	0	0	1	0	646	0.2%		
16	4/18/2020	7	0	0	0	0	518	0%		
17	4/25/2020	6	0	0	0	0	409	0%		


Influenza results by subtype, PCR testing only (NREVSS)





1

Public Health - Seattle & King County Summary of Influenza Syndromic and Laboratory Surveillance




King County syndromic surveillance

2015 - 16 2016-17 2**0**17 **-** 18 5 yr avg Percent 2018-19 2019-20 Baseline Current week 8 7 6 5 0-9 11 13 15 17 19 21 23 25 27 29 31 34 36 38 40 42 44 46 48 50 52 1 3 5 Week

Weekly Percent of ED Visits for ILI by Year

All ages Note: The change from ICD-9 to ICD-10 codes in October 2015 may impact trends. Last updated Apr 26, 2020 ; 'current week' is week ending Apr 25, 2020
Baseline: Mean % ILI during non-flu weeks for previous three seasons, adding two standard deviations A non-flu week is a period of 2+ consecutive weeks where each one accounted for <2% of the season's total number of specimens that tested positive for influenza by PCR.

Weekly Percent of ED Visits for Influenza-Like Illness By Age Group



ALLHOSPITALS, Last Updated

National data from CDC



A Weekly Influenza Surveillance Report Prepared by the Influenza Division Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet 2019-20 Influenza Season Week 16 ending Apr 18, 2020



does not, however, measure the extent of geographic spread of flu within a state. Therefore, outbreaks occurring in a single city could cause the state to display high activity levels.

picture of influenza activity for the whole state

picture or immeriza activity for the winderstate.

That displayed in this map are based on data collected in ILINet, whereas the State and Territorial flu activity map are based on reports from state and territorial epidemiologists. The data presented in this map is preliminary and may change as more data is received.

"Differences in the data presented by OCD and state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.

Please report any of the following:

- Laboratory-Confirmed influenza-associated deaths
- Patients with novel or unsubtypable influenza viruses
- · Outbreaks of influenza-like illness in a long-term care facility

Additional Resources:

Additional King County Flu Information, Resources, and Surveillance UW Virology Laboratory Respiratory Virus Surveillance Washington State Influenza Surveillance Update

Reporting Timeframe

Within 3 business days Immediately Immediately

National Influenza Update

Global Influenza Update

(206) 296-4774 Phone: (206) 296-4803 Fax:

Contact Information



Report updated on 4/29/2020

Public Health - Seattle & King County Summary of Influenza Deaths and Long-Term Care Facility (LTCF) Influenza Outbreaks

Confirme	ed cas	es as o	f wee	k 17 (endi	ng 04	/25/2	20)							
	2019-2020		2018	-2019	2017	7-2018	2016	5-2017	2015	2015-2016		2014-2015		ar avg	
Influenza Deaths in Week 17		0		6		0		0		0		1	1	.4	
Influenza deaths, season to date (since 9/29/2019)	33		4	19		41	8	82	:	16	4	1 1	45.8		
LTCF Outbreaks in Week 17	1			0	2			0		0		0	0	.4	
LTCF Outbreaks, season to date (since 9/29/2019)		25		43 65		65	92		17		61		55.6		
	201	2019-2020		2018-2019		2017-2018		2016-2017		2015-2016		2014-2015		5-year avg	
Total Seasonal LTCF Outbreaks		25		43		68		92		18		65		57.2	
Flu type:															
A	12	48%	37	86%	15	22%	62	67%	7	39%	49	75%	34	59%	
В	4	16%	0	0%	6	9%	3	3%	7	39%	4	6%	4	7%	
A and B	2	8%	1	2%	5	7%	4	4%	0	0%	2	3%	2.4	4%	
Info not available	7	28%	5	12%	42	62%	23	25%	4	22%	10	15%	16.8	29%	
			_												
	201	9-2020	2018	3-2019	2017-2018 2016-2017		2015-2016		2014-2015		5-year avg				
Total Seasonal Influenza Deaths	33			52	50		84		16		43		49		
Flu type:															
A	21	64%	48	92%	33	66%	75	89%	10	63%	40	93%	41.2	84%	
H1/H1N1	8	24%	11	21%	1	2%	1	1%	6	38%	0	0%	3.8	8%	
н3	1	3%	5	10%	6	12%	18	21%	1	6%	7	16%	7.4	15%	
A (not typed)	12	36%	32	62%	26	52%	56	67%	3	19%	33	77%	30	61%	
В	12	36%	2	4%	11	22%	7	8%	6	38%	3	7%	5.8	12%	
Not typed	0	0%	2	4%	6	12%	1	1%	0	0%	0	0%	1.8	4%	
Cour															
<u>Sex:</u> Male	18	55%	27	52%	17	34%	41	49%	7	44%	17	40%	21.8	44%	
Female	15	45%	25	48%	33	66%	43	51%	9	56%	26	60%	27.2	56%	
Terriale	13	43/0	23	4070	33	0070	43	31/0		3070	20	0070	27.2	3070	
Age:															
Under 5 years	1	3%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	
5 - 17	1	3%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	
18 - 44	4	12%	1	2%	0	0%	1	1%	3	19%	1	2%	1.2	2%	
45 - 64	4	12%	13	25%	7	14%	5	6%	5	31%	6	14%	7.2	15%	
65+ years	23	70%	38	73%	43	86%	78	93%	8	50%	36	84%	40.6	83%	
Average	66.1		73.6		81.1		81.9		64.9		81.7		76.6		
Race:															
White	19	58%	35	67%	33	66%	54	64%	12	75%	35	81%	33.8	69%	
Asian	2	6%	5	10%	2	4%	13	15%	2	13%	1	2%	4.6	9%	
Black	1	3%	1	2%	3	6%	4	5%	2	13%	5	12%	3	6%	
Amer Indian	1	3%	1	2%	0	0%	0	0%	0	0%	0	0%	0.2	0%	
Hispanic/Latino	3	9%	2	4%	2	4%	3	4%	0	0%	1	2%	1.6	3%	
Other	0	0%	0	0%	1	2%	1	1%	0	0%	1	2%	0.6	1%	
Unknown	7	21%	8	15%	9	18%	9	11%	0	0%	0	0%	5.2	11%	
Flu vaccine status															
Up to date	13	39%	16	31%	26	52%	39	46%	6	38%	21	49%	21.6	44%	
Not up to date	15	45%	19	37%	10	20%	20	24%	8	50%	5	12%	12.4	25%	
Unknown	5	15%	17	33%	14	28%	25	30%	2	13%	17	40%	15	31%	
CHAICWII	, J	13/0	17	J3/0	14	20/0	23	3070		rt upda				3	

3