Adapted from WA DOH Form 347-102

Fax Page 1 To: Public Health — Seattle & King County Confidential fax line: (206) 744 - 5622 CONFIDENTIAL SEXUALLY TRANSMITTED DISEASE CASE REPORT



Report STDs within three work days (WAC 246-101-101/301)

| PATIENT INFO | ORMATIO | N | | | | | | | | | | | | | |
|--|--------------|----------------------------------|----------------------|---|---|---|-----------|--------------------------------|--|---|--|--------|--------------|------------------|--|
| LAST NAME FI | | | | | FIRST NAME | | | MIDDLE NAME | | | DATE OF BIRTH | | | | |
| | | | | | | | | | | MO DAY | | YR | | | |
| ADDRESS | | | | | | CITY | | | | | STATE | | ZIP COD | DE | |
| | | | | | | | | | | | | | | | |
| TELEPHONE | | | EMAIL | | | | ENG | | 2 🗆 Vac | | DIAGNOS | | TE | | |
| | | | | | | | | | 、 | 1 | 1 1 | | | | |
| | | | | | | | | No (Lang) | | | MO DAY YR | | | | |
| SEX ASSIGNED AT BIRTH | | | | | | Transgender MTF | | | RACE (ch | CE (check all that apply) White | | | | | |
| | | □ Ma □ Fer | | | Transgender MTF Transgender FTM | | | | | | | | \Box Other | | |
| | | | nbinary/Gende | | Transgender FTM Other: Unknown | | | | American Indian / Alaskan Native | | | | | | |
| | | | | | | | | | | Native Hawaiian / Other Pacific Islander | | | | | |
| CURRENTLY REASON FOR EXA | | | AM | | R OF SEX PARTNERS (check all that apply) | | | HIV STATU | IV STATUS *Submit HIV/AIDS Case Report CURRENTLY | | | | | | |
| PREGNANT? (check one) | | | | Male | 6 | | | Previous positive ON PrEP? | | | | | | | |
| □ Yes □ Exposed □ No □ Symptor | | | nfection | Female | | | | | ew HIV diagnosis at this visit* | | | | | | |
| □ No □ Unknown | | tomatic ne Exam (No Symptoms) | | | □ Nonbinary / □ Ot | | | | | | egative HIV test at this visit | | | | |
| | 🗀 Routi | ne Exan | n (No Symptom | Genderqueer | nderqueer 🗌 Unknown | | | Did not test (unknown status) | | | | | | | |
| DIAGNOSIS - | DISFASE | | | | | | | | | | | | | | |
| | | | | | | | | | | | SYPHILIS | | | | |
| DIAGNOSIS (ch | | | SITES (all tha | t annly |): TREATMENT (d | heck al | pres | cribed): | | | check or | e): | | | |
| Asymptomatic | | | Cervix | c appiy | | Ceftriaxone: \Box 250 mg \Box 500 mg \Box 1 g | | | | | ary (Chan | | tc.) | | |
| Symptomati | icated | Urethra | | Cefixime: | | | | | | Secondary (Rash, etc.) | | | | | |
| Pelvic Inflammatory Disease Urin | | | | \square Azithromycin: \square 1 g \square 2 g | | | | | | Early Latent (< 1 year) | | | | | |
| 🗌 Ophthalmia | | | | Doxycycline: 100 mg BID x 7 days | | | | | | Unknown Duration or Late Latent | | | | | |
| Disseminated | | | | 🗌 Gentamicin: 🗌 240 mg | | | | | | Congenital | | | | | |
| □ Other Complications: □ Vagina □ Ocular | | | | | Gemifloxacin: 320 mg | | | | | | MANIFESTATIONS (check all that apply): | | | | |
| Othory | | | □ Ocular □ Other: | Other: | | | | | — D | □ Neurologic □ Otic □ Ocular □ Tertiary | | | | | |
| Date Tested: | | | | | Date Prescribe | a: | | | | | | | | | |
| CHLAMYDIA (I | | | | | •••••••• | | | /IENT (ch | | | | | | | |
| DIAGNOSIS (ch | - | | SITES (all that | | TREATMENT (check all prescribed): | | | | | Bicillin L - A: 🗌 2.4 MU IM x 1 | | | | | |
| Asymptoma | | icatod | Cervix | | □ Azithromycin: □ 1 g □ Doxycycline: □ 100 mg BID x 7 days | | | | | □ 2.4 MU IM x 3 | | | | | |
| Pelvic Inflam | | | | | \Box Levofloxacin: \Box 500 mg BiD x 7 days | | | | Doxycycline: 100 mg BID x 14 days | | | | | | |
| 🗌 Ophthalmia | | cuse | Rectum | | Other: | | | | 100 mg BID x 28 days | | | | | | |
| Other Comp | lications: | | Pharynx | | | | | | | Benzathine 50,000 units/kg IM x 1 | | | | | |
| | | | 🗌 Vagina | | | | | | PCN-G: 50,000 units/kg IM x 3 | | | | | | |
| | | | Ocular | | Data Duranih | I. | | | | Aqueous 18-24 MU/day IV Crystalline for 10.14 days | | | | | |
| Date Tested: Other: | | | Date Prescribed: | | | | | | for 10-14 days Penicillin G: | | | | | | |
| HERPES SIMPLEX DIAGNOSIS LABORATORY CONFIRMATIO | | | | | | OTHER DISEASES | | | | | | | | | |
| | | | Yes | | DN Chancroid | | | | Other: | | | | | | |
| | | | | | Lymphogranuloma Venereum | | | | Date Prescribed: | | | | | | |
| PARTNER TREATMENT PLAN (check one or more options) | | | | | | | | | | | | | | | |
| | | | - | | | son or b | y preso | ribing medicati | on for patier | nts to | give to th | eir se | x partne | rs (see side 2 | |
| Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2 for additional information). | | | | | | | | | | | | | | | |
| 🗌 In-person ev | aluation - N | umber o | of partners treat | ted follo | wing medical evaluation | ation: _ | | _ ! | Turn | over f | or Partner | reatm | ient Plan In | nstructions | |
| 🗌 Patient-deliv | vered treatm | nent* - N | Number of partr | ners for | whom provider pres | cribed of | or prov | vided expedited | partner the | rapy (| EPT) med | icatic | n pack to | be | |
| delivered by | the patient | to their | partner(s): | | *Patient-delivere | d treatm | nent is r | not recommende | d for men wl | no hav | ve sex with | n men | or patien | ts with syphilis | |
| REPORTING C | | ORMA | TION | | | | | | | | | | | | |
| DATE FACILITY NAME DIAGNOSING CLINICIAN | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | |
| ADDRESS | | | | | CITY | | | 5 | STATE | | Z | IP | | | |
| | | | | | | | | | | | | | | | |
| PERSON COMPLETING FORM | | | | | TELEPHON | TELEPHONE | | | EMAIL | MAIL | | | | | |
| | | | | | | () | | | | | | | | | |
| | | | | | () | | | | | | | | | | |
| | ٦ | Thank y | ou for reportin | ig an ST | () D. All information | will be | mana | ged with the st | rictest conf | ident | iality. | | | | |

use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile. Page 1

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

King County recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication, the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia, and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Disease Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call Public Health - Seattle & King County: (206) 744-3590.

Other STDs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM

**For persons weight >150 kg (~300 lbs), the ceftriaxone dose should be 1g IM

Alternatives for uncomplicated urethral, vaginal or rectal infection:

Azithromycin 2g PO as a single dose PLUS Gentamicin 240mg IM as a single dose

OR

Cefixime 800 mg PO as a single dose

**NOTE: There is no recommended alternative therapy for pharyngeal gonorrhea. Contact a local infectious disease or STD expert for assistance.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days (preferred) OR

Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR Erythromycin Ethylsuccinate 800 mg PO QID for 7 days **OR** Ofloxacin 300 mg PO BID for 7 days OR Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<u>http://www.cdc.gov/std/tg2015/default.htm</u>) for further information on treating pregnant patients, infections of the pharynx, treatment of infants and other details.

DOH 347-102, updated 03/08/2021. For persons with disabilities or persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).